



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

GROUP THEORY.

3. Proposed by L. E. DICKSON, Ph. D., The University of Chicago.

Show that the equation $x^4 - ax^3 + bx^2 - ax + 1 = 0$, where a and b are rational numbers, while neither $2 - b + (\frac{1}{2}a)^2$ nor $(1 + \frac{1}{2}b)^2 - a^2$ is the square of a rational number, is irreducible in the domain R of rational numbers; determine its group for this domain.

Solution by R. L. Börger, A. B., Graduate Student, The University of Chicago.

If reducible it could assume either of the forms :

$$(1) \quad (x^2 + px + 1)(x^2 + qx + 1) = x^4 + (p+q)x^3 + (2+pq)x^2 + (p+q)x + 1 = 0.$$

$$(2) \quad (x^2 + px + r)(x^2 + qx + \frac{1}{r}) = x^4 + (p+q)x^3 + (r + \frac{1}{r} + pq)x^2 + (qr + \frac{p}{r})x + 1 = 0$$

From (1), equating coefficients,

$$p+q = -a; \quad pq = b-2. \quad \therefore \left. \begin{matrix} p \\ q \end{matrix} \right\} = -\frac{1}{2}a \pm \sqrt{(\frac{1}{2}a)^2 + 2 - b}.$$

$$\text{From (2), } p+q = -a; \quad qr + \frac{p}{r} = -a; \quad r + \frac{1}{r} + pq = b. \quad \therefore p = -\frac{ar}{r+1}; \quad q = -\frac{a}{r+1}.$$

Substituting these in last we get

$$\frac{r^2+1}{r} + a^2 \frac{r}{(r+1)^2} = b, \text{ or } \frac{(r+1)^2}{r} + a^2 \frac{r}{(r+1)^2} = b+2. \quad \text{Put } \frac{(r+1)^2}{r} = x.$$

Then we have,

$$x^2 - (b+2)x + a^2 = 0, \text{ or } x = \frac{b+2}{2} \pm \sqrt{\frac{(b+2)^2}{4} - a^2}.$$

Now if the radicals for the above two quadratics are irrational then the original equation can not be reducible. Hence the equation is irreducible if neither $(2-b) + (\frac{1}{2}a)^2$ nor $(1 + \frac{1}{2}b)^2 - a^2$ is the square of a rational number.

To determine the group of the equation. Call its roots $\alpha_0, \beta_0, \alpha_1, \beta_1$ where the β 's are reciprocals of the α 's. Then $\alpha_0\beta_0 + \alpha_1\beta_1 = 2$. Its conjugates under G_4 are $\alpha_1\beta_0 + \alpha_0\beta_1$ and $\alpha_0\alpha_1 + \beta_0\beta_1$. No two of them can be equal for then the equation would have equal roots, which is impossible because of its irreducibility. The group is then either either G_8 consisting of

$$[1, (\alpha_0\beta_0); (\alpha_1\beta_1); (\alpha_0\beta_0)(\alpha_1\beta_1); (\alpha_0\beta_1)(\alpha_1\beta_0); (\alpha_0\alpha_1)(\beta_0\beta_1); (\alpha_0\alpha_1\beta_0\beta_1); (\alpha_1\alpha_0\beta_1\beta_0)], \text{ or}$$

$$(a) \ 1; (\alpha_0\beta_0); (\alpha_1\beta_1); (\alpha_0\beta_0)(\alpha_1\beta_1),$$

$$(b) \ 1; (\alpha_0\beta_0)(\alpha_1\beta_1); (\alpha_0\alpha_1\beta_0\beta_1); (\alpha_1\alpha_0\beta_1\beta_0),$$

$$(c) \ 1; (\alpha_0\beta_0)(\alpha_1\beta_1); (\alpha_0\alpha_1)(\beta_0\beta_1); (\alpha_0\beta_1)(\alpha_1\beta_0).$$

The group must be transitive since the equation is irreducible. Hence we may exclude (a).

It cannot be (b) since (b) is the regular cyclic group, and the equation, if this were its group, would be Abelian.

We compute $(\alpha_0 - \beta_0)(\alpha_1 - \beta_1)$ which belongs to the group (c):

$$\alpha_0\beta_0 + \alpha_1\beta_1 + \alpha_0\beta_1 + \alpha_1\beta_0 + \alpha_0\alpha_1 = b.$$

$$\therefore \alpha_0\beta_1 + \alpha_1\beta_0 + \alpha_0\alpha_1 + \beta_0\beta_1 = b - 2.$$

Factoring this, $(\alpha_0 + \beta_0)(\beta_1 + \beta_1) = b - 2$. Squaring, and remembering that $\alpha_0\beta_0 = \alpha_1\beta_1 = 1$,

$$(\alpha_0 - \beta_0)^2(\alpha_1 - \beta_1)^2 = [(\alpha_0 + \beta_0)^2 - 4][(\alpha_1 + \beta_1)^2 - 4].$$

$\therefore (\alpha_0 - \beta_0)(\alpha_1 - \beta_1) = 2\sqrt{[(1 + \frac{1}{2}b)^2 - a^2]}$, and as the irrationality of the radical is one of the conditions for irreducibility this group cannot be the group of the equation. The group of the equation is therefore G_8 .

PROBLEMS FOR SOLUTION.

ALGEBRA.

208. Proposed by F. P. MATZ, Sc. D., Ph. D., Professor of Mathematics and Astronomy in Defiance College Defiance, O.

$$\text{Solve } x^4 + y^4 = 14x^2y^2; \ x + y = m.$$

209. Proposed by F. P. MATZ, Sc. D., Ph. D., Professor of Mathematics and Astronomy in Defiance College, Defiance, O.

$$\text{Prove that } (a^4 + b^4 + c^4 + d^4) > 4abcd.$$

210. Proposed by W. J. GREENSTREET, A. M., Editor of The Mathematical Gazette, Stroud, England.

The sum of five quantities and the sum of their cubes are both zero. Show that the sum of their fifth powers is a factor of the sum of any odd powers of the quantities.

GEOMETRY.

236. Proposed by J. R. HITT.

If two sides of a triangle pass through a fixed point, the third side touches a fixed circle.